

Antidiabetic Activities of Triterpenoids Isolated from Bitter Melon Associated with Activation of the AMPK Pathway

Min-Jia Tan, Ji-Ming Ye, Nigel Turner, Cordula Hohnen-Behrens, Chang-Qiang Ke, Chun-Ping Tang, Tong Chen, Hans-Christoph Weiss, Ernst-Rudolf Gesing, Alex Rowland, David E. James,* and Yang Ye*

*Correspondence: d.james@garvan.org.au (D.E.J.), yye@mail.shcnc.ac.cn (Y.Y.)

DOI 10.1016/j.chembiol.2008.05.003

(Chemistry & Biology 15, 263–273, March 2008)

The ^1H NMR assignments of atomic numbers from 25 to 30 in Table 1 were inadvertently mistyped.

They should be corrected as follows:

Table 1. ^{13}C and ^1H NMR Data of Compounds 2, 3, 5–7 in $\text{C}_5\text{D}_5\text{N}$

	δ_{H}				
	2	3	5	6	7
25					
26	1.65 s	1.72 s	1.81 s	1.82 s	1.70 s
27	1.67 s	1.73 s	1.70 s	1.71 s	1.68 s
28	1.56 s	1.45 s	1.43 s	1.44 s	1.53 s
29	0.87 s	0.86 s	1.00 s	1.03 s	1.09 s
30	0.93 s	0.83 s	0.78 s	0.78 s	0.83 s